

4-2016

Effects of distillers' dried grains with solubles and soybean oil on dietary lipid, fiber, and amino acid digestibility in corn-based diets fed to growing pigs

N. A. Gutierrez
Iowa State University

Nick V. L. Serão
Iowa State University, serao@iastate.edu

John F. Patience
Iowa State University, jfp@iastate.edu

Follow this and additional works at: https://lib.dr.iastate.edu/ans_pubs



Part of the [Agriculture Commons](#), [Animal Experimentation and Research Commons](#), and the [Animal Sciences Commons](#)

The complete bibliographic information for this item can be found at https://lib.dr.iastate.edu/ans_pubs/814. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Animal Science at Iowa State University Digital Repository. It has been accepted for inclusion in Animal Science Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Effects of distillers' dried grains with solubles and soybean oil on dietary lipid, fiber, and amino acid digestibility in corn-based diets fed to growing pigs

Abstract

The use of corn co-products increases the concentration of fiber and often the use of supplemental lipids in swine diets, which may affect energy and nutrient digestibility. An experiment was conducted to determine the effects of reduced oil distillers dried grains with solubles (DDGS) and soybean oil (SBO) on dietary AA, acid hydrolyzed ether extract (AEE), and NDF digestibility in corn-based diets fed to growing pigs. Eighteen growing pigs (BW = 33.8 ± 2.2 kg) were surgically fitted with a T-cannula in the distal ileum and allocated to 1 of 6 dietary treatment groups in a 3-period incomplete Latin square design, with 9 observations per treatment. Six dietary treatments were obtained by adding 0, 20, and 40% DDGS to corn-casein diets formulated with 2 and 6% SBO. Ileal digesta and fecal samples were collected and the apparent ileal (AID) and total tract digestibility (ATTD) of AEE and NDF, and the AID of Lys were determined. Apparent values were corrected for endogenous losses of lipids and true ileal (TID) and true total tract digestibility (TTTD) values were calculated. Results showed that the AID of Lys decreased ($P < 0.001$) with the inclusion of DDGS, but was not affected ($P = 0.63$) by the inclusion of SBO. An interaction between DDGS and SBO on the AID ($P = 0.002$) and ATTD ($P = 0.009$) of NDF was observed, where the AID and ATTD of NDF decreased with DDGS at 6% SBO, but no effect was observed at 2% SBO. The AID of NDF increased with SBO at 0% DDGS, but no effect was observed at 20 or 40% DDGS. An interaction between DDGS and SBO on the AID ($P = 0.011$) and ATTD ($P = 0.008$) of AEE was observed, where the AID and ATTD of AEE increased with SBO. The AID and ATTD of AEE increased with DDGS at 2% SBO, but no effect was observed at 6% SBO. Correction by ileal and fecal endogenous loss of AEE (9.5 and 13.6 g/kg of DMI) showed that increasing dietary AEE had no effect on the TID and TTD of AEE ($P > 0.05$). In conclusion, the AID of Lys decreased with DDGS and was not affected by lipids from SBO. Greatest AID and ATTD of NDF was observed in diets with a high AEE and low NDF content. Low values of apparent digestibility of AEE in lower lipid diets are possibly the result of endogenous losses of lipids, because the true digestibility of AEE was not affected by the dietary increase of AEE.

Keywords

Distillers dried grains with solubles, soybean oil, fiber, lipid, pigs

Disciplines

Agriculture | Animal Experimentation and Research | Animal Sciences

Comments

This is a manuscript of an article published as Gutierrez, N. A., N. V. L. Serão, and J. F. Patience. "Effects of distillers' dried grains with solubles and soybean oil on dietary lipid, fiber, and amino acid digestibility in corn-based diets fed to growing pigs." *Journal of animal science* 94, no. 4 (2016): 1508-1519. doi:[10.2527/jas.2015-9529](https://doi.org/10.2527/jas.2015-9529). Posted with permission.

RUNNING HEAD: DDGS and soybean oil on digestibility

**Effects of distillers dried grains with solubles and soybean oil on dietary lipid, fiber, and
amino acid digestibility in corn based diets fed to growing pigs**

N. A. Gutierrez, N. V. L. Serão, and J. F. Patience¹

Department of Animal Science, Iowa State University, Ames, IA 50011

¹ Corresponding author: 201B Kildee Hall, Ames, IA 50011; jfp@iastate.edu

ABSTRACT

The use of corn co-products increases the concentration of fiber and often the use of supplemental lipids in swine diets, which may affect energy and nutrient digestibility. An experiment was conducted to determine the effects of reduced oil distillers dried grains with solubles (**DDGS**) and soybean oil (**SBO**) on dietary AA, acid hydrolyzed ether extract (**AEE**), and NDF digestibility in corn-based diets fed to growing pigs. Eighteen growing pigs (BW = 33.8 ± 2.2 kg) were surgically fitted with a T-cannula in the distal ileum and allocated to 1 of 6 dietary treatment groups in a 3-period incomplete Latin square design, with 9 observations per treatment. Six dietary treatments were obtained by adding 0, 20, and 40% DDGS to corn-casein diets formulated with 2 and 6% SBO. Ileal digesta and fecal samples were collected and the apparent ileal (**AID**) and total tract digestibility (**ATTD**) of AEE and NDF, and the AID of Lys were determined. Apparent values were corrected for endogenous losses of lipids and true ileal (**TID**) and true total tract digestibility (**TTTD**) values were calculated. Results showed that the AID of Lys decreased ($P < 0.001$) with the inclusion of DDGS, but was not affected ($P = 0.63$) by the inclusion of SBO. An interaction between DDGS and SBO on the AID ($P = 0.002$) and ATTD ($P = 0.009$) of NDF was observed, where the AID and ATTD of NDF decreased with DDGS at 6% SBO, but no effect was observed at 2% SBO. The AID of NDF increased with SBO at 0% DDGS, but no effect was observed at 20 or 40% DDGS. An interaction between DDGS and SBO on the AID ($P = 0.011$) and ATTD ($P = 0.008$) of AEE was observed, where the AID and ATTD of AEE increased with SBO. The AID and ATTD of AEE increased with DDGS at 2% SBO, but no effect was observed at 6% SBO. Correction by ileal and fecal endogenous loss of AEE (9.5 and 13.6 g/kg of DMI) showed that increasing dietary AEE had no effect on the TID and TTD of AEE ($P > 0.05$). In conclusion, the AID of Lys decreased with

42 DDGS and was not affected by lipids from SBO. Greatest AID and ATTD of NDF was observed
43 in diets with a high AEE and low NDF content. Low values of apparent digestibility of AEE in
44 lower lipid diets are possibly the result of endogenous losses of lipids, because the true
45 digestibility of AEE was not affected by the dietary increase of AEE.

46 **Keywords:** Distillers dried grains with solubles, soybean oil, fiber, lipid, pigs

47

48

50 Corn co-products are rich in dietary fiber and their inclusion in swine diets dilutes the
51 traditionally starch-based dietary energy (Kerr et al, 2013). Dietary fiber from corn and its co-
52 products is only partially fermented in the gastrointestinal tract of pigs because it is rich in
53 insoluble non-starch polysaccharides, such as cellulose, arabinoxylans, and lignin (Bach
54 Knudsen, 1997; Jaworski, 2012). Substitution of highly digestible nutrients and the limited
55 fermentability of fiber in diets including corn co-products result in a linear decline of dietary
56 energy digestibility (Noblet and Perez, 1993; Le Goff and Noblet, 2001; van Milgen, 2006).
57 Technology advances in the ethanol industry have decreased the starch and lipid concentrations
58 in distillers dried grains with solubles (**DDGS**), concentrating fiber in corn co-products, and
59 resulting in the necessity to add greater quantities of lipids to the diet to maintain an acceptable
60 concentration of dietary energy. Lipids in swine diets typically originate from intact lipids in
61 feed ingredient matrices or extracted lipids such as soybean oil (**SBO**) or any number of
62 vegetable oils (Azain, 2001; Adams and Jensen, 1984).

63 Extracted lipid sources induce smaller amounts of endogenous lipid losses and have
64 greater digestibility than intact lipid sources when fed to growing pigs (Kil et al., 2010; Kim et
65 al., 2013). Additionally, the inclusion of DDGS in swine diets decrease the apparent digestibility
66 of dietary fiber, AA, and energy in swine diets (Urriola and Stein, 2010; Gutierrez et al., 2013).
67 However, the effect of varying amounts of extracted lipids added to diets containing DDGS on
68 the digestibility of fiber, lipids, and AA has not been investigated.

The objective of the present study was therefore to establish the relationships between the dietary increase in fiber and lipids from DDGS and SBO on the digestibility of AA, NDF, and acid hydrolyzed ether extract (AEE) in growing pigs..

MATERIALS AND METHODS

The experimental protocol was reviewed and approved by the Institutional Animal Care and Use Committee at Iowa State University (12-13-7686-S).

Animals, Housing, and Experimental Design

Eighteen growing barrows (progeny of sire line 337 × dam line C-22, PIC, Hendersonville, TN) were housed in individual pens (1.2 × 1.2 m) equipped with a feeder, a cup waterer, and a half-slatted concrete floor in an environmentally controlled building. All pigs were surgically fitted with a T-cannula in the distal ileum following procedures described by Stein et al. (1998). After recovery from surgery, pigs were weighed (initial BW = 33.8 ± 2.2 kg) and randomly allocated to 1 of 6 dietary treatments groups in a 3-period incomplete Latin square design, resulting in 9 observations per treatment. Pigs were not allowed to repeat dietary treatments across periods. Each collection period involved 9 d of adaptation to dietary treatments followed by 2 d of feces sub-sample collection and 3 d of ileal digesta sub-sample collection.

Dietary Treatments

Ingredient composition of the dietary treatments is shown in Table 1. Six dietary treatments were obtained by adding 0, 20, and 40% DDGS to corn-casein diets formulated with 2% and 6% SBO. The same batch of DDGS was used for all dietary treatments and values of

nutrient composition were obtained from NRC (2012). A corn-casein diet with 2% SBO was formulated to meet or exceed the nutrient requirements of growing pigs (NRC, 2012), and 5 additional dietary treatments were obtained by adding DDGS and SBO at the expense of corn. Casein and the portion of the diet including limestone, monocalcium phosphate, salt, vitamins and trace mineral premixes were maintained constant. Diets also contained 0.5% Cr₂O₃ as an inert marker. All pigs received the same daily amount of feed, which was provided at a level of approximately 90% of predicted *ad-libitum* intake of the diet formulated with 2% SBO and no DDGS. Ad-libitum intake was estimated as DE intake (kcal/d) = $13.162 \times (1 - e^{-0.0176 \times BW})$, where BW was average BW (NRC, 1987). The daily feed allowance was divided into 2 equal meals provided at 0700 and 1600 h. At the end of each collection period, all pigs were weighed and daily feed allowance for the next collection period was adjusted.

Sample Collection

After 9 d of adaptation to the diet, feces were collected via grab sampling on d 10 and 11, and stored at -20°C. On d 12, 13, and 14, ileal digesta samples were collected for 8 h by attaching a 207-mL plastic bag (Whirl-Pak, Nasco, Fort Atkinson, WI) to the opened cannula with a cable tie. Bags were removed whenever they were filled with digesta or at least every 30 min, and stored at -20°C to prevent bacterial degradation. At the conclusion of each experimental period, frozen ileal and fecal samples were allowed to thaw at room temperature and pooled within animal, with a sub-sample collected for chemical analysis. Ileal sub-samples were lyophilized before chemical analysis. Fecal sub-samples were oven-dried in a convection oven at 65°C to constant weight (Jacobs et al., 2011). Samples of diets were collected weekly, throughout the experimental period, and in the feed mill during mixing. At the conclusion of the experiment, dietary samples were pooled and sub-sampled for chemical analysis. Diets, feed

ingredients, and dried ileal and fecal sub-samples were finely ground in a Wiley Mill (Variable Speed Digital ED-5 Wiley Mill; Thomas Scientific, Swedesboro, NJ) through a 1-mm screen and stored in desiccators to maintain a constant percentage of DM.

Chemical Analysis and Calculations

Samples of diets were analyzed for DM (method 930.15; AOAC Int., 2007), GE by a bomb calorimeter (Model 6200; Parr Instrument Co., Moline, IL) with benzoic acid as a standard (6,318 kcal GE/kg of benzoic acid; Parr Instrument Co., Moline, IL), acid hydrolyzed ether extract (**AEE**; Sanderson, 1986; Soxtec 2050, FOSS North America, Eden Prairie, MN), starch (method 996.11; AOAC Int., 2007), ADF (Goering and Van Soest, 1970), neutral detergent fiber (NDF; Van Soest and Robertson, 1980), total dietary fiber (**TDF**; method 985.29; AOAC Int., 2007), and N using the combustion method (method 990.03: AOAC International, 2007) with a Trumac apparatus (Leco Corporation, St. Joseph, MI) with EDTA for calibration ($9.58 \pm 0.01\%$ N; Leco Corporation, St. Joseph, MO). Crude protein was calculated as nitrogen \times 6.25. Ileal digesta and fecal samples were also analyzed for DM, GE, NDF, and AEE. Diets and ileal digesta were analyzed for AA (University of Missouri Agriculture Experiment Station Chemical Laboratories, Columbia, MO) according to method 982.30 E (a, b, c; AOAC Int., 2007). Chromium was determined in diets, ileal digesta, and fecal sub-samples using the method of Fenton and Fenton (1979) and absorption was measured at 440 nm using a spectrophotometer (Synergy 4, BioTek, Winooski, VT). Chromic oxide standard samples were assayed to confirm the accuracy of the analytical procedure, and a recovery of $100.8 \pm 1.95\%$ was attained.

The DE value was determined by multiplying the GE by the observed ATTD of GE of the ingredient, and the ME was estimated from the calculated DE and CP of the ingredient

(Noblet and Perez, 1993). For each dietary treatment, the AID and ATTD of DM, GE, NDF, and AEE, and the AID of AA were calculated (Oresanya et al., 2008). True ileal digestibility (**TID**) and true total tract digestibility (**TTTD**) of AEE were calculated using the regression method (Jørgensen et al., 1993). Following the procedures outlined by Kil et al. (2010), the total apparently digested AEE (g/kg of DMI) at the end of the ileum (**TAD_i**) and over the entire intestinal tract (**TAD_t**) were estimated by multiplying the AID and ATTD of AEE by the dietary AEE con (g/kg of DM) of each pig, respectively. A linear response of both the TAD_i and TAD_t of AEE to the increase in dietary AEE intake (g/kg of DM) was expected, and the Y-intercepts of these two linear functions were considered to represent the estimated endogenous losses of AEE (g/kg of DMI) at the end of the ileum and over the entire intestinal tract, respectively (Jørgensen et al., 1993; Kil et al., 2010). The slopes of the linear functions are considered the average true digestibility of AEE at the ileal and total tract level. The TID and TTTD of AEE in diets were also calculated by correcting individual values of AID and ATTD for endogenous losses of AEE (g/kg of DMI) at the end of the ileum and over the entire intestinal tract, respectively (Stein et al., 2007; Kil et al., 2010).

Statistical Analyses

Univariate Analysis and Normality Test. The data were analyzed in a mixed model (PROC MIXED, SAS Inst., Inc., Cary, NC) including the fixed effects of DDGS, SBO, their interaction and the covariate initial BW, and the random effects of Period (3 levels) and Group (6 levels with 3 pigs per group).

Studentized residuals from each analysis were used to test normality. Outliers were removed until the Shapiro-Wilk's test reached $P > 0.05$ and Studentized residuals fell within $\pm 3\sigma$.

Least squares means for DDGS and interaction between DDGS and SBO were compared using Tukey-Kramer adjustment. Effects and multiple comparison differences were deemed significant at $\alpha \leq 0.05$.

Response Surface Model. Orthogonal polynomial contrasts were used to test the linear and quadratic effects of DDGS and SBO, when possible. Linear effects tested were: DDGS (L_DDGS), SBO (L_SBO), and interaction L_SBO*L_DDGS. Quadratic effects tested were: Q_DDGS and the interaction L_SBO*Q_DDGS.

Depending on the significance ($P \leq 0.05$) of the contrasts, response surface models were generated to predict response according to the dietary concentrations of DDGS and SBO. A hierarchical model selection was used to construct the response surface models. Models were only constructed when both DDGS and SBO were significant ($P \leq 0.05$), regardless of the orthogonal polynomial effect (linear or quadratic). If the quadratic effect of Q_DDGS was significant ($P \leq 0.05$), the linear effect of L_DDGS was kept in the model. Similarly, if the interaction L_SBO*Q_DDGS was significant ($P \leq 0.05$), the interaction L_SBO*L_DDGS was kept in the model, as well as all other effects following the hierarchy of the model (L_SBO, L_DDGS, and Q_DDGS), since these effects are part of the interaction effect of L_SBO*Q_DDGS. Since SBO had only two levels (2 and 6%), we could only test the linear effect (L_SBO and interactions).

Estimation of endogenous losses of lipids. In addition to the analyses of the raw data for AID and ATTD of AEE, these traits were adjusted for the estimated endogenous losses of AEE and reanalyzed as TID and TTTD. Estimation of the endogenous loss of AEE for each lipid trait was performed using a linear regression of the ileal and fecal TAD AEE on the AEE intake from

the 6 dietary treatments. The 6 dietary treatments represent the combination of the 2 SBO and 3 DDGS. The intercept of the ileal and fecal models was considered the endogenous loss of AEE at each compartment, respectively. The TID and TTTD of AEE were then estimated as:

$$Y_{ij} = \frac{(TAD_{ij} - \text{endogenous loss of AEE})}{\text{intake of AEE}_j}$$

where: Y_{ij} is the estimated TID or TTTD of AEE of the i^{th} observation in the j^{th} dietary treatment ($j=1$ to 6), TAD_{ij} is the observed value of TAD_i or TAD_t AEE, and intake of AEE_j is the intake of AEE of the j^{th} dietary treatment. The estimated TID and TTTD of AEE were analyzed using the same mixed model previously described. All statistical analyses were performed using SAS 9.3 (Statistical Analysis Software, SAS Institute Inc., Cary, NC, USA).

RESULTS AND DISCUSSION

All pigs successfully recovered from surgery, remained healthy and readily consumed their diets throughout the experiment. The NDF and AEE of the DDGS used in the present study were 39.2% and 7.1%, respectively. The analyzed nutrient composition of diets showed that dietary inclusion of DDGS increased the dietary AEE in treatments formulated with 2% SBO (4.4, 5.7, and 6.3% AEE) and 6% SBO (8.4, 9.1, and 10.1% AEE: Table 2). The dietary NDF also increased with DDGS in treatments formulated with 2% SBO (6.9, 11.0, and 14.6% NDF) and 6% SBO (6.8, 9.7, and 14% NDF).

Apparent Digestibility of Energy and Amino Acids

The AID and ATTD of GE and DM decreased with dietary inclusion of DDGS ($P < 0.001$; Table 3), resulting in DE and ME values of diets formulated with 2% and 6% SBO decreasing ($P < 0.01$) with DDGS addition (Table 4). The decrease in dietary energy is a direct result of gradually replacing highly digestible starch with low-digestible dietary fiber from DDGS. In growing pigs it has been previously estimated that % increase in dietary NDF reduced the energy digestibility by 0.9% (Le Goff and Noblet, 2001). Similar effects on apparent digestibility of DM have been previously reported after insoluble dietary fiber was included at the expense of a highly digestible source of carbohydrates (Schulze et al., 1994; Wilfart et al., 2007). In contrast, the dietary inclusion of SBO increased the AID of GE ($P = 0.046$), resulting in DE and ME values of diets increasing ($P < 0.01$) by approximately 0.2 Mcal/kg of ME when SBO increased from 2% to 6% in the diet. The large effect of adding DDGS and SBO on the apparent digestibility of DM, GE, and on DE and ME values is supported by the response surface models (Table 5), where a high predictability of AID of GE ($R^2 = 0.78$), DM ($R^2 = 0.86$), DE ($R^2 = 0.83$), and ME ($R^2 = 0.87$) from SBO and DDGS inclusion was observed.

The AID of all indispensable AA decreased ($P < 0.05$) with DDGS and, with the exception of the AID of Ile ($P = 0.015$) and Met ($P = 0.008$), were not affected by SBO ($P > 0.05$). A decline in the AID of AA from the dietary increase of co-products from the corn-ethanol distillation industry has also been previously reported (Urriola and Stein, 2010; Gutierrez et al., 2013). The decrease in AID of AA observed may result from the combination of an increase in the amount of endogenous N loss and a decrease in absorption of endogenous and exogenous N. Dietary fiber intake has been associated with the increase of endogenous loss of nitrogen (Souffrant, 1991, 2001; Moughan, 2003), and insoluble fiber in particular associated with the secretion of pancreatic juice (Langlois et al., 1987). Although the increase of dietary

fiber may reduce the digestibility of AA (Schulze et al., 1994), insoluble dietary fiber, such as corn fiber, has been reported to have only minor effects on the digestibility of dietary AA (Zhu et al., 2005) and on the basal endogenous losses of AA (Leterme et al., 1996). Therefore, the formation of biologically unavailable Maillard reactions products from excessive heating, and the addition of solubles, is likely the cause of the lower AID of Lys in DDGS (Pahm et al., 2008; Stein and Shurson, 2009).

Apparent Digestibility of Fiber

An interaction between DDGS and SBO on the AID ($P = 0.002$) and ATTD ($P = 0.009$) of NDF was observed. The dietary increase of DDGS at 2% SBO, and the subsequent increase in dietary insoluble NDF, showed no effect on the apparent digestibility of NDF at the terminal ileum and over the entire intestinal tract (Figure 1). Similar results on the AID of insoluble dietary fiber have been previously reported (Graham et al., 1986; Urriola and Stein, 2010; Gutierrez et al., 2013), and AID of NDF values observed in the present study ($\approx 28\%$ AID of NDF at 2% SBO) agree with previously reported data from corn bran (Gutierrez et al., 2013). Fermentation of dietary fiber at the terminal ileum is well documented, and the extent varies with the solubility of dietary fiber (Jørgensen et al., 1996; Noblet and Le Goff, 2001; Urriola and Stein, 2010; Gutierrez et al., 2013). Surprisingly, the ATTD of NDF was not affected by DDGS increase in diets with 2% SBO, which differ from previous reports where the increase in insoluble dietary fiber from corn origin decreased the ATTD of dietary fiber (Urriola and Stein, 2010; Gutierrez et al., 2013).

Conversely, the dietary increase of DDGS in diets with 6% SBO, and the subsequent increase in dietary insoluble NDF, resulted in a reduction of the ileal and fecal apparent digestibility of NDF. The different response to DDGS addition between diets with 2% and 6%

SBO is possibly the consequence of the elevated dietary lipid concentration in the latter set of diets, and the ability of dietary lipids to increase the retention time of digesta in low fiber diets. Dietary lipids have been previously reported to increase the retention time of digesta in pigs (Cervantes-Pahm and Stein, 2008) and in laying hens (Mateos et al., 1982). An increase in the retention time of digesta may also increase the fermentability of fiber due to a longer period of exposure of substrates to the intestinal microbiota at the terminal ileum or in the hindgut (Morel et al., 2006; Wilfart et al., 2007). This effect may be noticeable at 0% DDGS and 6% SBO, where dietary NDF is low but AEE is high, and where the AID and ATTD of NDF were the greatest (46.4% and 52.0%, respectively). It is also evident when the AID and ATTD of NDF increased with the increase of SBO in diets with the lowest fiber content (0% DDGS).

The possibility of a longer transit time in pigs fed diets with 6% SBO may be offset by the dietary increase of NDF from DDGS, decreasing in turn the apparent digestibility of NDF. A dietary increase of NDF from DDGS may stimulate bowel movement and reduce the transit time of digesta (Bastianelli et al., 1996; Schneeman, 1998; Bindelle et al., 2008), which agree with data from pigs fed insoluble dietary fiber from wheat bran where a decrease in retention time of digesta was reported (Wilfart et al., 2007). The moderate predictability of AID and ATTD of NDF ($R^2 = 0.53$ and $R^2 = 0.36$, respectively) from SBO and DDGS may provide evidence the difficulty of the interaction between a fibrous and lipid-rich ingredients.

Values of apparent digestibility of NDF were greater over the entire intestinal tract than at the terminal ileum. This observation suggests that part of fiber from corn origin is fermented at the terminal ileum and an additional portion is fermented in the hindgut of pigs (Urriola and Stein, 2010; Gutierrez et al., 2013). The value of ATTD for dietary fiber is therefore an accurate estimate of the total fermentation of dietary fiber in pigs.

Apparent Digestibility of Acid Hydrolyzed Ether Extract

The dietary AEE increased with SBO and DDGS inclusion. An interaction between DDGS and SBO on the AID ($P = 0.011$) and ATTD ($P = 0.008$) of AEE was observed, where the AID and ATTD of AEE increased with the dietary level of AEE from SBO, and increased likewise with the dietary level of AEE from DDGS but only when added to 2% SBO diets (Table 6). An interaction between DDGS and SBO on the TAD_i ($P = 0.001$) and TAD_t ($P = 0.001$) was also observed, where the TAD_i and TAD_t increased with DDGS at the two levels of SBO, but values were greater at 6% SBO than at 2% SBO. A similar interaction between dietary lipid and fiber concentrations on the apparent digestibility of dietary lipids has also been previously reported (Dégen et al., 2009), and the positive relationship between dietary increase and apparent digestibility of AEE observed in the present study agree with previously reported data from pigs (Just et al., 1980; Jørgensen et al., 1993; Kim et al., 2013).

The observed response in apparent digestibility of AEE was modulated by the source of dietary lipids. This was evidenced by the response surface models, where the magnitude of the slopes for the effects of SBO on AID and ATTD of AEE were greater than those of DDGS. This observation is supported by the high predictability of the AID and ATTD of AEE ($R^2 = 0.68$ and $R^2 = 0.79$, respectively) from DDGS and SBO inclusion (Figure 2). The shape of the observed response originate from the fact that apparent digestibility of extracted lipid is greater than for intact lipids, as previously reported in pigs fed diets supplemented with SBO (Agunbiade et al., 1992), palm kernel oil (Agunbiade et al., 1999), sunflower oil (San Juan and Villamide, 2000), and corn oil (Kim et al., 2013). Therefore, greater values of AID and ATTD of AEE were observed in diets formulated with 6% SBO, where most of the AEE content is extracted lipids, than in diets formulated with 2% SBO, where most of the AEE content is intact lipids. Values of

AID and ATTD of AEE in the present study fall within the range of values reported by Kil et al. (2010) for diets formulated with extracted or intact lipids.

However, low apparent digestibility values for AEE from diets formulated with low levels of dietary AEE may result from the contribution of endogenous losses of AEE to the total AEE output (Jørgensen et al., 1993; Kil et al., 2010; Kim et al., 2013). Similar effects of endogenous losses on the apparent digestibility of dietary AA from have also been reported previously (Fan and Sauer, 1997).

The increase of dietary fiber intake may stimulate the synthesis of endogenous microbial lipids in the hindgut, increasing in turn the endogenous losses of lipids at the total tract level and reducing the ATTD of lipids. A net synthesis of lipids in the hindgut of pigs has been previously reported (Shi and Noblet, 1993; Bakker, 1996). Kil et al. (2010) reported greater values of AID vs. ATTD of AEE from semi-purified diets formulated with increasing levels of fiber from Solka-Floc. These results suggest that dietary lipids are digested and absorbed before the end of the ileum, and that differences between values of AID and ATTD of AEE are the product of endogenous microbial lipid synthesis in the hindgut of pigs (Low, 1980; Drackley, 2000). Corn and DDGS were the only fiber sources in the present study. Fiber from corn origin is fermented partially because of its chemical composition; a small portion is fermented by microbes at the ileal level, and the majority of corn dietary fiber reaches the hindgut where is moderately fermented (Gutierrez et al., 2013). The increase in corn fiber reaching the hindgut may stimulate microbial activity and lipid synthesis (Eyssen, 1973; Bach Knudsen et al., 1991), increasing endogenous losses of lipids from microbial origin, which may explain the lower values in ATTD of AEE relative to AID observed in the present study. Values of AID of AEE therefore reflect more accurately the digestibility of dietary AEE than ATTD values.

In the present study, dietary NDF increased similarly with inclusion of DDGS in diets containing 2% or 6% SBO. A decrease in the apparent digestibility of lipids with a dietary increase of NDF has been previously reported (Just et al., 1980; Bakker, 1996; Hansen et al., 2006), but the reduction has been associated with characteristics of dietary fiber such as solubility and viscosity, rather than its dietary concentration (Fahey et al., 1990; Bach Knudsen and Hansen, 1991; Smits and Annison, 1996). Fiber in corn and its co-products is highly insoluble and has low viscosity (Jaworski, 2012; Gutierrez et al., 2014); therefore, the increase of dietary fiber from the dietary inclusion of DDGS may not negatively affect the apparent digestibility of AEE at 2% or 6% SBO. This assumption is supported by Kil et al. (2010), who observed no effects of dietary increase of highly insoluble and low viscous purified NDF on the apparent digestibility of AEE.

Endogenous Losses and True Digestibility of Acid Hydrolyzed Ether Extract

The TAD_i and TAD_t of AEE increased ($P < 0.01$) with AEE intake (Figure 3), and the estimated linear functions explained 99 % of the variation in TAD_i and TAD_t. Linearity is a prerequisite of the linear regression method, and the observed linear response supports its use for the estimation of endogenous loss of AEE from both compartments of the digestive system (Jørgensen et al., 1993; Fan and Sauer, 1997). The endogenous losses of AEE, corresponding to the y-intercepts of the linear functions, were 9.47 g/kg of DMI at the end of the ileum and 13.64 g/kg of DMI over the entire intestinal tract. The average true digestibility of AEE at the end of the ileum and over the entire intestinal tract, estimated from the slope of the linear functions, was 94 % and 93 %, respectively. The TID and TTTD of AEE (Table 7), calculated by adjusting the AID and ATTD for endogenous losses, showed that increasing the dietary inclusion of SBO or DDGS had no effect ($P > 0.05$) on TID or TTTD.

The source of dietary lipids may result in different estimates of endogenous lipid loss at both the ileal and total tract levels. Estimates of endogenous losses of AEE from feeding intact lipids reported by Kil et al. (2010) were greater than estimates from extracted lipids, at both the ileal and total tract levels. Greater endogenous loss of AEE from feeding intact lipids may result from the commonly greater concentrations of fiber in ingredients containing intact lipids. Fiber in the feed ingredient may depress absorption of dietary lipids and resorption of endogenous lipids, resulting in greater endogenous lipid loss (Bach Knudsen and Hansen, 1991). Dietary treatments in the present study were formulated with varying ratios of intact to extracted lipids; therefore, estimates of endogenous losses of AEE for these diets are not directly comparable to estimates reported previously for intact or extracted lipid sources. Additionally, discrepancies between estimates of endogenous lipid loss may result from the fact that the range in dietary lipid concentrations differed among experiments. Although pigs in the present study were not offered diets with very low AEE levels, the regression curves calculated were able to explain approximately 99% of the variation in totally apparently digested AEE. Other factors such as analytical method to estimate lipid concentrations (ether extract vs. AEE), lipid sources (Kim et al., 2013), non-lipid dietary components (Jørgensen et al., 1992b), and animal effects (Jørgensen et al., 2012a) may also affect the estimates of endogenous lipid loss.

Greater true digestibility of extracted lipids compared with intact lipids at both collection sites has been reported previously (Kil et al., 2010). This observation results from the fact that intact lipids are encased within cell membranes, therefore are more resistant to the formation of emulsions and enzymatic digestion than extracted sources (Adams and Jensen, 1984; Bach Knudsen et al., 1993). The average TID (94.2%) and TTTD (93.1%) of AEE estimated in this experiment are similar to values for extracted lipids (93.8 and 94.2%, respectively) reported by

Kil et al. (2010). The average TTTD value also falls within the range of values (91.2 to 97.7%) reported for the TTTD of SBO (Adams and Jensen, 1984; Jørgensen et al., 1993; Jørgensen and Fernández, 2000). The similarity between estimates of true digestibility in the present study and previous estimates for extracted lipids may rely on the fact that SBO was an important contributor to the total lipid content in experimental diets, ranging from approximately 30% to 67%. Additionally, an important portion of the lipids in DDGS has been previously extracted from the corn grain and added back in the solubles prior to drying (Díaz-Royón et al., 2012). Its digestion should therefore be similar to an extracted source. As outlined by Kil et al. (2010), additional variation in TID and TTD for AEE among experiments can be introduced because of differences in dietary lipid intake, diet composition, diet processing, and lipid sources.

Although the apparent digestibility of AEE increased with the inclusion of dietary AEE from SBO and DDGS, the true digestibility of AEE was not affected by the dietary concentration of AEE. The reason for this observation is that the endogenous loss of AEE affects the apparent digestibility, but not the true digestibility of AEE. These results therefore support the hypothesis that the true digestibility of a lipid source is not affected by the amount of dietary lipid intake (Freeman et al., 1968; Adams and Jensen, 1984).

Conclusions

Results from the present experiment showed that the AID of AA was negatively associated with the concentration of DDGS. No effect on the AID and ATTD of NDF from the dietary increase of NDF from DDGS was observed in diets with low concentrations of AEE from SBO. However, in diets with high concentrations of AEE from SBO, and low NDF from DDGS, the AID and ATTD of NDF was greatest. The high concentration of AEE, and low concentration

of NDF, may increase the retention time of digesta and therefore the fermentability of NDF in the gastrointestinal tract. The AID and ATTD of AEE increased with dietary AEE inclusion from SBO and DDGS. Low values of apparent digestibility of AEE in diets with low concentrations of AEE are possibly the result of endogenous losses of AEE, because the true digestibility of AEE was not affected by the dietary increase of AEE.

LITERATURE CITED

- Adams, K. L., and A. H. Jensen. 1984. Comparative utilization of in-seed fats and the respective extracted fats by the young pig. *J. Anim. Sci.* 59:1557-1566.
- Adams, K. L., and A. H. Jensen. 1985. Effect of dietary protein and fat levels on the utilization of the fat in sunflower seeds by the young pig. *Anim. Feed Sci. Technol.* 13:159-170.
- Agunbiade, J. A., J. Wiseman, and D. J. A. Cole. 1992. Utilization of dietary energy and nutrients from soybean products by growing pigs. *Anim. Feed Sci. Technol.* 36:303-318.
- Agunbiade, J. A., J. Wiseman, and D. J. A. Cole. 1999. Energy and nutrient use of palm kernels, palm kernel meal and palm kernel oil in diets for growing pigs. *Anim. Feed Sci. Technol.* 80:165-181.
- AOAC International. 2007. Official methods of analysis of AOAC Int. 18th ed. Assoc. Off. Anal. Chem. Int., Gaithersburg, MD.
- Azain, M. J. 2001. Fat in swine nutrition. Page 95-105. In *Swine Nutrition*. 2nd ed. A. J. Lewis and L. L. Southern, ed. CRC Press New York, NY.
- Bach Knudsen, K. E. 1997. Carbohydrate and lignin contents of plant materials used in animal feeding. *Anim. Feed Sci. Technol.* 67:319-338.

403 Bach Knudsen, K. E., and I. Hansen. 1991. Gastrointestinal implications in pigs of wheat and oat
 404 fractions. 1. Digestibility and bulking properties of polysaccharides and other major
 405 constituents. *Br. J. Nutr.* 65:217-232.

406 Bach Knudsen, K. E., B. B. Jensen, J. O. Andersen, and I. Hansen. 1991. Gastrointestinal
 407 implications in pigs of wheat and oat fractions. 2. Microbial activity in the
 408 gastrointestinal tract. *Br. J. Nutr.* 65:233-248.

409 Bach Knudsen, K. E., B. B. Jensen, and I. Hansen. 1993. Digestion of polysaccharides and other
 410 major components in the small and large intestine of pigs fed on diets consisting of oat
 411 fractions rich in β -D-glucan. *Br. J. Nutr.* 70:537-556.

412 Bakker, G. C. M. 1996. Interaction between carbohydrates and fat in pigs: impact on energy
 413 evaluation of feeds. PhD Diss. Wageningen Agric. Univ., Wageningen, The Netherlands.

414 Bastianelli, D., D. Sauvant, and A. Rerat. 1996. Mathematical modeling of digestion and nutrient
 415 absorption in the pig. *J. Anim. Sci.* 74:1873-1887.

416 Bindelle, J., P. Leterme, and A. Buldgen. 2008. Nutritional and environmental consequences of
 417 dietary fibre in pig nutrition: A review. *Biotechnol. Agron. Soc. Environ.* 12:69-80.

418 Cervantes-Pahm, S. K. and H. H. Stein. 2008. Effects of soybean oil and soybean protein
 419 concentration on the concentration of digestible amino acids in soybean products fed to
 420 growing pigs. *J. Anim. Sci.* 86:1841-1849.

421 Dégen, L., V. Halas, J. Tossenberger, Cs. Szabó, and L. Babinszky. 2009. The impact of dietary
 422 fiber and fat levels on total tract digestibility on energy and nutrients in growing pigs and
 423 its consequence for diet formulation. *Acta Agric. Scand. Sect. A. Anim. Sci.* 59:150-160.

424 Díaz-Royón, F., A. García, and K. A. Rosentrater. 2012. Composition of fat in distillers grains.
 425 Report. iGrow. South Dakota State University Extension Publication.

http://lib.dr.iastate.edu/cgi/viewcontent.cgi?article=1673&context=abe_eng_pubs

(Accessed 29 June 2015).

Drackley, J. K. 2000. Lipid metabolism. Page 97-119 in *Farm animal Metabolism and Nutrition*.

J. P. F. D;Mello, ed. CAB Int. Publ., Wallingford, Oxfordshire, UK.

Eyssen, H. 1973. Role of the gut microflora in the metabolism of lipids and sterols. *Proc. Nutr.*

Soc. 32:59-63.

Fahey, G. C. Jr., N. R. Merchen, J. E. Corbin, A. K. Hamilton, K. A. Serbe, and D. A. Hirakawa.

1990. Dietary fiber for dogs: II. Iso-total dietary fiber (TDF) additions of divergent fiber

sources to dog diets and their effects on nutrient intake, digestibility, metabolizable

energy and digesta mean retention time. *J. Anim. Sci.* 68:4229-4235.

Fan, M. Z. and W. C. Sauer. 1997. Determination of true ileal amino acid digestibility in

feedstuffs for pigs with the linear relationships between distal ileal outputs and dietary

inputs of amino acids. *J. Sci. Food Agric.* 73:189-199.

Fenton, T. W. and M. Fenton. 1979. An improved procedure for the determination of chromic

oxide in feed and feces. *Can. J. Anim. Sci.* 59: 631-634.

Freeman, C. P., D. W. Holme, and E. F. Annison. 1968. The determination of the true

digestibilities of interesterified fats in young pigs. *Br. J. Nutr.* 22:651-660.

Goering, H. K. and P. J. Van Soest. 1970. Forage fiber analyses (apparatus, reagents, procedures,

and some applications). In: *Handbook No. 379. USDA-ARS, Washington, DC.*

Graham, H., K. Hesselman, and P. Åman 1986. The influence of wheat bran and sugar-beet pulp

on the digestibility of dietary components in a cereal-based pig diet. *J. Nutr.* 116:242-

251.

448 Gutierrez, N. A., B. J. Kerr, and J. F. Patience. 2013. Effect of insoluble-low fermentable fiber
 449 from corn-ethanol distillation origin on energy, fiber, and amino acid digestibility,
 450 hindgut degradability of fiber, and growth performance of pigs. J. Anim. Sci. 91:5314-
 451 5325.

452 Gutierrez, N. A., N. V. L. Seroo, B. J. Kerr, R. T. Zijlstra, and J. F. Patience. 2014. Relationships
 453 among dietary fiber components and the digestibility of energy, dietary fiber, and amino
 454 acids and energy content of nine corn coproducts fed to growing pigs. J. Anim. Sci.
 455 92:4505-4517.

456 Farrell, D. J. 1973. Digestibility by pigs of the major chemical components of diets high in plant
 457 cell-wall constituents. Anim. Prod. 16:43-47.

458 Jacobs, B. M., J. F. Patience, W. A. Dozier III, K. J. Stalder and B. J. Kerr. 2011. Effects of
 459 drying methods on nitrogen and energy concentrations in pig feces and urine, and poultry
 460 excreta. J. Anim. Sci. 89:2624-2630.

461 Jaworski, N. 2012. Carbohydrate composition, *in vitro* digestion, and effects of xylanase and
 462 phytase on nutrient and energy digestibility by pigs in grains and grain coproducts. M.S.
 463 Thesis. Univ. of Illinois, Urbana-Champaign, IL.

464 Jørgensen, H., and J. A. Fernández. 2000. Chemical composition and energy value of different
 465 fat sources for growing pigs. Acta Agric. Scand. Sect. A. Anim. Sci. 50:129-136.

466 Jørgensen, H., K. Jakobsen, and B. O. Eggum. 1992a. Excretion of biliary fat and fatty acids in
 467 growing pigs. J. Anim. Feed Sci. 1:139-149.

468 Jørgensen, H., K. Jakobsen, and B. O. Eggum. 1992b. The influence of different protein, fat, and
 469 mineral levels on the digestibility of fat and fatty acids measured at the terminal ileum
 470 and in faeces of growing pigs. Acta Agric. Scand. Sect. A. Anim. Sci. 42:177-184.

471 Jørgensen, H., K. Jakobsen, and B. O. Eggum. 1993. Determination of endogenous fat and fatty
 472 acids at the terminal ileum and on faeces in growing pigs. *Acta Agric. Scand. Sect. A.*
 473 *Anim. Sci.* 42:177-184.

474 Jørgensen, H., X. Q. Zaho, and B. O. Eggum. 1996. The influence of dietary fibre and
 475 environmental temperature on the development of the gastrointestinal tract, digestibility,
 476 degree of fermentation in the hindgut, and energy metabolism in pigs. *Br. J. Nutr.*
 477 75:365-378.

478 Just, A., J. O. Andersen, and H. Jørgensen. 1980. The influence of diet composition on the
 479 apparent digestibility of crude fat and fatty acids at the terminal ileum and overall in pigs.
 480 *Z. Tierphysiol. Tierernaehr. Futtermittelkde.* 44:82-92.

481 Kerr, B. J., W. A. Dozier, and G. C. Shurson. 2013. Effects of reduced-oil corn distillers dried
 482 grains with solubles composition on digestible and metabolizable energy value and
 483 prediction in growing pigs. *J. Anim. Sci.* 91:3231-3243.

484 Kil, D. J., T. E. Sauber, D. B. Jones, and H. H. Stein. 2010. Effect of the form of dietary fat and
 485 the concentration of dietary neutral detergent fiber on ileal and total tract endogenous
 486 losses and apparent and true digestibility of fat by growing pigs. *J. Anim. Sci.* 88:2959-
 487 2967.

488 Kim, B. G., D. Y. Kil, and H. H. Stein. 2013. In growing pigs, the true ileal and total tract
 489 digestibility of acid hydrolyzed ether extract in extracted corn oil is greater than in intact
 490 sources of corn oil or soybean oil. *J. Anim. Sci.* 91:755-763.

491 Langlois, A., T. Corring, and C. Février. 1987. Effect of wheat bran on exocrine pancreas
 492 secretion in the pig. *Reprod. Nutr. Dév.* 27:929-939.

493 Le Goff, G., and J. Noblet. 2001. Comparative digestibility of dietary energy and nutrients in
 494 growing pigs and adult sows. *J. Anim. Sci.* 79:2418-2427.

495 Leterme, P., P. van Leeuwen, A. Thewis, and J. Huisman. 1996. Chemical composition of pea
 496 inner fiber isolates and their effect on endogenous digestive secretions in pigs. *J. Sci.*
 497 *Food Agric.* 72:127-134.

498 Low, A. G. 1980. Nutrient absorption in pigs. *J. Sci. Food Agric.* 31:1087-1130.

499 Mateos, G. G., J. L. Sell, and J. A. Eastwood. 1982. Rate of food passage (transit time) as
 500 influenced by level of supplemental fat. *Poult. Sci.* 61:94-100.

501 Morel, P. C. H., T. S. Lee, and P. J. Moughan. 2006. Effect of feeding level, live weight and
 502 genotype on the apparent faecal digestibility of energy and organic matter in the growing
 503 pig. *Anim. Feed Sci. Technol.* 126:63-74.

504 Moughan, P. J. 2003. Amino acid digestibility and availability in food and feedstuffs. In: R. O.
 505 Ball, editor, *Digestive Physiology in pigs. Proc. 9th Intl. Symp. Vol. 1.* Univ. Alberta,
 506 Alberta, Canada. p. 199-221.

507 Noblet, J. and J. M. Perez. 1993. Prediction of digestibility of nutrients and energy values of pig
 508 diets from chemical analysis. *J. Anim. Sci.* 71:3389-3398.

509 Noblet, J. and G. Le Goff. 2001. Effect of dietary fibre on the energy value of feeds for pigs.
 510 *Anim. Feed Sci. Technol.* 90:35-52.

511 NRC. 1987. Predicting feed intake of food-producing animals. *Natl. Acad. Press, Washington,*
 512 *D.C.*

513 NRC. 2012. *Nutrient Requirements of Swine. 11th rev. ed.* *Natl. Acad. Press, Washington, DC.*

514 Oresanya, T.F., A.D. Beaulieu and J.F. Patience. 2008. Investigations of energy metabolism in
 515 weanling barrows: The interaction of dietary energy concentration and daily feed
 516 (energy) intake. *J. Anim. Sci.* 86: 348-363.

517 Pahm, A. A., C. Pedersen, and H. H. Stein. 2008. Application of the reactive lysine procedure to
 518 estimate lysine digestibility in distillers dried grains with solubles fed to growing pigs. *J.*
 519 *Agric. Food Chem.* 56:9441–9446.

520 San Juan, L. D., and M. J. Villamide. 2000. Nutritional evaluation of sunflower seed and
 521 products derived from them. Effect of oil extraction. *Br. Poult. Sci.* 41:182-192.

522 Sanderson, P. 1986. A new method of analysis of feedingstuffs for the determination of crude
 523 oils and fats. In W. Haresign and D. J. A. Cole (ed.) *Recent Advances in Animal*
 524 *Nutrition.* Butterworths, London, UK. p. 77-81.

525 Schneeman, B. O. 1998. Dietary fiber and gastrointestinal function. *Nutr. Res.* 4:625-632.

526 Schulze, H., P. van Leeuwen, M. W. A. Verstegen, J. Huisman, W. B. Souffrant, and F. Ahrens.
 527 1994. Effect of level of dietary neutral detergent fiber on ileal apparent digestibility and
 528 ileal nitrogen losses in pigs. *J. Anim. Sci.* 72:2362-2368.

529 Souffrant, W. B. 1991. Endogenous nitrogen losses during digestion in pigs. In: M. W. A.
 530 Verstegen, J. Huisman, and L. A. den Hartog, editors, *Digestive physiology in pigs.*
 531 *EAAP Publ. No. 54.* Pudoc, Wageningen, The Netherlands. p. 147.

532 Souffrant, W. B. 2001. Effect of dietary fiber on ileal digestibility and endogenous nitrogen
 533 losses in the pig. *Anim. Feed Sci. Technol.* 90:93-102.

534 Shi, X. S., and J. Noblet. 1993. Contribution of the hindgut to digestion of diets in growing pigs
 535 and adult sows: Effect of diet composition. *Livest. Prod. Sci.* 34:237-252.

536 Smits, C. H. M., and G. Annison. 1996. Non-starch plant polysaccharides in broiler nutrition –
 537 Towards a physiologically valid approach to their determination. *World's Poult. Sci. J.*
 538 52:204-221.

539 Stein, H. H., C. F. Shipley, and R. A. Easter. 1998. Technical Note: A technique for inserting a
 540 T-cannula into the distal ileum of pregnant sows. *J. Anim. Sci.* 76:1433-1436.

541 Stein, H. H., M. L. Gibson, C. Pedersen, and M. G. Boersma. 2006. Amino acid and energy
 542 digestibility in ten samples of distillers dried grain with solubles fed to growing pigs. *J.*
 543 *Anim. Sci.* 84:853-860.

544 Stein, H. H., B. Sève, M. F. Fuller, P. J. Moughan, and C. F. M. de Lange. 2007. Invited review:
 545 Amino acid bioavailability and digestibility in pig feed ingredients: Terminology and
 546 application. *J. Anim. Sci.* 85:172-180.

547 Urriola, P. E., D. Hoehler, C. Pedersen, H. H. Stein, and G. C. Shurson. 2009. Amino acid
 548 digestibility of distillers dried grains with solubles, produced from sorghum, a sorghum-
 549 corn blend, and corn fed to growing pigs. *J. Anim. Sci.* 87:2574-2580.

550 Stein, H. H., and G. C. Shurson. 2009. Board invited review. The use and application of distillers
 551 dried grains with solubles in swine diets. *J. Anim. Sci.* 87:1292-1303.

552 Urriola, P. E., G. C. Shurson, and H. H. Stein. 2010. Digestibility of dietary fiber in distillers
 553 coproducts fed to growing pigs. *J. Anim. Sci.* 88:2373-2381.

554 Urriola, P. E. and H. H. Stein. 2010. Effects of distillers dried grains with solubles on amino
 555 acid, energy, and fiber digestibility and hindgut fermentation of dietary fiber in a corn-
 556 soybean meal diet fed to growing pigs. *J. Anim. Sci.* 88:1454-1462.

557 Van Milgen, J. 2006. Nutrient flow models, energy transactions and energy feed systems. In: R.
 558 M. Gous, T. R. Morris, and C. Fisher, editors, Mechanistic modeling in pig and poultry
 559 production. CABI Publishing, Wallingford, Oxfordshire, UK. p. 143-145.
 560 Van Soest, P. J. and J. B. Robertson. 1980. Systems of analysis for evaluating fibrous feeds. In:
 561 Workshop Standardization of Analytical Methodology for Feeds, Int. Dev. Res Center,
 562 Ottawa, ON, Canada. Pp. 49-60.
 563 Wilfart, A., L. Montagne, P. H. Simmins, J. Noblet, and J. van Milgen. 2007. Digesta transit in
 564 different segments of the gastrointestinal tract of pigs as affected by insoluble fibre
 565 supplied by wheat bran. Br. J. Nutr. 98:54-62.
 566 Zhu, C. L., M. Rademacher, and C. F. M. de Lange. 2005. Increasing dietary pectin level reduces
 567 utilization of digestible threonine intake, but not lysine intake, for body protein
 568 deposition in growing pigs. J. Anim. Sci. 83:1044-1053.
 569

570 **Table 1.** Ingredient composition (%) of the experimental diets (as-fed basis)¹

Item	SBO, %	2			6		
	DDGS, %	0	20	40	0	20	40
Ingredient, %							
Corn		81.7	61.7	41.7	77.7	57.7	37.7
DDGS		0.0	20.0	40.0	0.0	20.0	40.0
Casein		12.5	12.5	12.5	12.5	12.5	12.5
Soybean oil		2.0	2.0	2.0	6.0	6.0	6.0
Limestone		1.3	1.3	1.3	1.3	1.3	1.3
Monocalcium phosphate		1.0	1.0	1.0	1.0	1.0	1.0
Chromic oxide		0.5	0.5	0.5	0.5	0.5	0.5
Vitamin premix ²		0.2	0.2	0.2	0.2	0.2	0.2
Trace mineral premix ³		0.2	0.2	0.2	0.2	0.2	0.2
Salt		0.6	0.6	0.6	0.6	0.6	0.6
Energy and nutrients ⁴							
ME, Mcal/kg		3.39	3.33	3.27	3.59	3.54	3.48
SID Lys, %		0.98	1.03	1.08	0.98	1.02	1.07
SID Thr, %		0.61	0.71	0.80	0.61	0.70	0.80
SID Met+Cys, %		0.60	0.70	0.80	0.59	0.69	0.78
SID Trp, %		0.20	0.22	0.23	0.20	0.21	0.23

571 ¹SBO = soybean oil; DDGS = reduced oil distillers dried grains with solubles.

572 ²Provided per kilogram of complete diet: 6,614 IU of vitamin A; 827 IU of vitamin D; 26 IU
573 of vitamin E; 2.6 mg of vitamin K; 29.8 mg of niacin; 16.5 mg of pantothenic acid; 5.0 mg of

574 riboflavin; 0.023 mg of vitamin B₁₂.

575 ³Provided per kilogram of complete diet: Zn, 165 mg as ZnSO₄; Fe, 165 mg as FeSO₄; Mn, 39

576 mg as MnSO₄; Cu, 17 mg as CuSO₄; I, 0.3 mg as Ca(IO₃)₂; and Se, 0.3 mg as Na₂SeO₃.

577 ⁴Values were calculated (NRC, 2012). SID = standardized ileal digestible.

578 **Table 2.** Analyzed nutrient composition of the experimental diets (% as-fed basis)¹

Item	SBO, %	2			6			DDGS	SBO
	DDGS, %	0	20	40	0	20	40		
DM, %		89.1	88.0	89.5	89.2	89.8	89.5	89.8	-
GE, Mcal/kg		3.98	4.08	4.18	4.18	4.33	4.42	4.40	9.16
AEE, %		4.4	5.7	6.3	8.4	9.1	10.1	7.1	98.1
Starch, %		51.1	39.2	27.2	48.6	36.7	24.7	2.9	-
ADF, %		2.4	3.4	4.6	2.2	3.4	4.5	15.8	-
NDF, %		6.9	11.0	14.6	6.8	9.7	14.0	33.8	-
CP		17.9	21.8	25.7	17.5	21.4	25.4	26.7	-
Indispensable AA									
Arg		0.7	0.8	1.0	0.6	0.8	1.0	1.3	-
His		0.5	0.6	0.8	0.5	0.6	0.8	0.9	-
Ile		0.9	1.0	1.2	0.9	1.0	1.2	1.1	-
Leu		2.0	2.3	2.8	1.9	2.3	2.9	3.3	-
Lys		1.2	1.3	1.4	1.1	1.2	1.5	1.0	-
Met		0.5	0.5	0.6	0.5	0.5	0.6	0.6	-
Phe		1.0	1.1	1.3	0.9	1.1	1.3	1.4	-
Thr		0.8	0.9	1.0	0.7	0.9	1.1	1.1	-
Trp		0.2	0.2	0.3	0.2	0.2	0.3	0.2	-
Val		1.1	1.3	1.5	1.1	1.2	1.5	1.4	-
Dispensable AA ²									
Mean		10.1	11.6	13.3	11.2	13.5	10.1	1.1	-

All AA ³	18.9	21.7	25.1	21.0	25.6	18.9	1.2	-
---------------------	------	------	------	------	------	------	-----	---

579 ¹SBO = soybean oil; DDGS = distillers dried grains with solubles; AEE = acid hydrolyzed

580 ether extract; TDF = total dietary fiber.

581 ²Sum of Ala, Asp, Cys, Glu, Gly, Pro, Ser, and Tyr.

582 ³Sum of all indispensable and dispensable AA.

583 **Table 3.** Apparent ileal digestibility (AID) and apparent total tract digestibility (ATTD) of components of experimental diets^{1,2}

SBO, %		2			6			P-value			
Item	DDGS, %	0	20	40	0	20	40	pSEM	SBO	DDGS	SBO x DDGS
AID, %											
GE		80.6 ^a	73.3 ^{cb}	70.0 ^c	84.0 ^a	74.3 ^b	70.1 ^c	0.96	0.046	<0.001	0.178
DM		80.2 ^a	71.1 ^b	66.1 ^c	83.4 ^a	71.4 ^b	65.0 ^c	0.96	0.325	<0.001	0.076
NDF		32.5 ^b	26.8 ^b	25.3 ^b	46.4 ^a	21.7 ^b	22.4 ^b	3.12	0.389	<0.001	0.002
ATTD, %											
GE		87.8 ^a	84.0 ^b	80.0 ^c	88.3 ^a	84.3 ^b	80.0 ^c	0.48	0.509	<0.001	0.813
DM		88.3 ^a	83.8 ^b	79.0 ^c	88.7 ^a	83.9 ^b	78.3 ^c	0.46	0.892	<0.001	0.424
NDF		47.3 ^{ab}	48.7 ^{ab}	45.5 ^{bc}	52.0 ^a	45.7 ^{abc}	40.9 ^c	2.28	0.451	<0.001	0.009
AID of AA, %											
Arg		84.2 ^{ab}	81.9 ^b	83.2 ^b	87.0 ^a	82.4 ^b	83.2 ^{ab}	1.06	0.061	0.001	0.145
His		87.1 ^a	82.8 ^b	83.1 ^b	89.5 ^a	82.6 ^b	83.1 ^b	0.99	0.160	<0.001	0.107
Ile		87.6 ^a	82.5 ^b	81.8 ^b	88.9 ^a	82.9 ^b	83.3 ^b	0.67	0.015	<0.001	0.539
Leu		90.1 ^a	87.0 ^b	87.5 ^b	91.3 ^a	86.9 ^b	87.7 ^b	0.73	0.239	<0.001	0.435

Lys	88.9 ^a	85.2 ^b	84.6 ^b	90.2 ^a	84.3 ^b	84.7 ^b	0.82	0.629	<0.001	0.127
Met	92.8 ^b	90.0 ^c	89.6 ^c	93.7 ^a	90.1 ^c	90.1 ^c	0.44	0.008	<0.001	0.223
Phe	90.1 ^a	86.9 ^b	87.3 ^b	91.2 ^a	86.6 ^b	87.3 ^b	0.79	0.588	<0.001	0.390
Thr	81.4 ^a	75.4 ^b	74.6 ^b	82.8 ^a	74.7 ^b	75.6 ^b	1.18	0.371	<0.001	0.434
Trp	89.2 ^a	84.6 ^b	84.4 ^b	90.7 ^a	83.7 ^b	84.4 ^b	1.04	0.754	<0.001	0.242
Val	86.7 ^a	81.7 ^b	81.7 ^b	87.9 ^a	81.8 ^b	82.4 ^b	0.86	0.191	<0.001	0.665
Indispensable ³	88.1 ^a	84.0 ^b	84.1 ^b	89.4 ^a	83.9 ^b	84.4 ^b	0.81	0.253	<0.001	0.388
Dispensable ⁴	84.9 ^a	80.9 ^b	80.0 ^b	87.5 ^a	81.0 ^b	80.8 ^b	1.02	0.052	<0.001	0.256
All AA ⁵	86.4 ^a	82.4 ^b	82.0 ^b	88.4 ^a	82.4 ^b	82.5 ^b	0.88	0.096	<0.001	0.274

pSEM = Pooled SEM

^{a,b,c,d} Means within a row lacking a common superscript letter are different ($P \leq 0.05$).

¹Least squares means of 9 pigs per diet.

²SBO = soybean oil; DDGS = reduced oil distillers dried grains with solubles; AEE = acid hydrolyzed ether extract.

588 ³Average AID for all indispensable AA.

589 ⁴Average AID for all dispensable AA.

590 ⁵Average AID for all AA (indispensable and dispensable).

591

592 **Table 4.** Estimated digestible and metabolizable energy value of diets^{1,2}

SBO, %		2			6			pSEM	<i>P</i> -value		
Item	DDGS-RO, %	0	20	40	0	20	40		SBO	DDGS-RO	SBO*DDGS-RO
As-fed basis, Mcal/kg											
	DE	3.49 ^{cb}	3.43 ^c	3.34 ^d	3.69 ^a	3.65 ^a	3.53 ^b	0.020	<0.001	<0.001	0.639
	ME	3.37 ^b	3.28 ^c	3.17 ^d	3.57 ^a	3.49 ^a	3.35 ^{cb}	0.019	<0.001	<0.001	0.617

593 pSEM = Pooled SEM

594 ^{a,b,c}Means within a row lacking a common superscript letter are different ($P \leq 0.05$).

595 ¹ Least squares means of 9 pigs per diet.

596 ²SBO = soybean oil; DDGS-RO = reduced oil distillers dried grains with solubles.

597

598 **Table 5.** Regression coefficients of the response surface models for the effects of soybean oil (SBO) and reduced oil distillers dried
599 grains with solubles (DDGS) on apparent digestibility and energy value of diets^{1,2}

Trait	Regression components						R ²
	Intercept	SBO	DDGS	DDGS ²	SBO*DDGS	SBO*DDGS ²	
AID							
GE	81.2908	0.3100	-0.5447	0.0057	-	-	0.78
		(0.046)	(<0.001)	(0.005)	(0.071)	(0.651)	
DM	78.9176	0.7571	-0.5133	0.0056	-0.0318	-	0.86
		(0.325)	(<0.001)	(0.005)	(0.026)	(0.629)	
NDF	24.4572	3.7063	0.5498	-0.0118	-0.3960	0.0069	0.53
		(0.389)	(<0.001)	(0.005)	(0.003)	(0.037)	
AEE	67.2575	2.7325	0.2469	-5*10 ⁻⁵	-0.0425	-	0.68
		(<0.001)	(0.001)	(0.894)	(0.003)	(0.800)	
ATTD							
NDF	44.8490	1.2009	0.2205	-0.0026	-0.0722	-	0.36
		(0.451)	(<0.001)	(0.571)	(0.004)	(0.256)	

AEE	55.3292	3.5898 (<0.001)	0.3654 (<0.001)	-0.0002 (0.413)	-0.0512 (0.004)	- (0.153)	0.79
AID of AA							
Met	93.0344	0.0949 (0.008)	-0.2458 (<0.001)	0.0039 (<0.001)	- (0.442)	- (0.111)	0.71
Energy concentration, as-fed basis							
DE	3.4098	0.0483 (<0.001)	-0.0041 (<0.001)	- (0.123)	- (0.745)	- (0.384)	0.83
ME	3.2922	0.0469 (<0.001)	-0.0053 (<0.001)	- (0.133)	- (0.670)	- (0.387)	0.87

¹Traits included were simultaneously significant for SBO and DDGS (linear or quadratic) or any of the possible interactions.

²*P*-values in brackets.

Table 6. Apparent digestibility and total apparently digested acid hydrolyzed ether extract (AEE) at the end of the ileum and over the entire intestinal tract of diets containing extracted or intact lipids^{1,2}

		SBO, %			2			6				
		DDGS, %			0	20	40	0	20	40	<i>P</i> -value	
Item	AEE intake, g/kg of DM	44.2	57.3	63.3	84.2	90.9	100.9	pSEM	SBO	DDGS	SBO*DDGS	
AID of AEE, %		72.5 ^c	75.9 ^{bc}	79.1 ^b	83.5 ^a	83.4 ^a	83.9 ^a	1.05	<0.001	0.005	0.011	
TAD _i AEE, g/kg of DMI		32.0 ^f	43.3 ^e	49.7 ^d	70.4 ^c	75.6 ^b	84.4 ^a	0.72	<0.001	<0.001	0.001	
ATTD of AEE, %		62.6 ^d	67.6 ^c	71.6 ^b	77.5 ^a	77.0 ^a	80.4 ^a	1.30	<0.001	<0.001	0.008	
TAD _t AEE, g/kg of DMI		27.8 ^f	38.5 ^e	45.0 ^d	65.2 ^c	69.9 ^b	80.9 ^a	0.89	<0.001	<0.001	0.001	

pSEM = Pooled SEM

^{a,b,c}Means within a row lacking a common superscript letter are different ($P \leq 0.05$).

¹ Least squares means of 9 pigs per diet.

²SBO = soybean oil; DDGS = reduced oil distillers dried grains with solubles; AID = apparent ileal digestibility; ATTD = apparent total tract digestibility. TAD_i = total apparently digested at the end of the ileum; TAD_t = total apparently digested over the entire intestinal tract.

Table 7. True ileal digestibility (TID) and true total tract digestibility (TTTD) of acid hydrolyzed ether extract (AEE) of diets containing extracted or intact lipids^{1,2}

Item	SBO, %	2			6			pSEM	<i>P</i> -value		
	DDGS, %	0	20	40	0	20	40		SBO	DDGS	SBO*DDGS
TID of AEE, %		94.0	92.5	94.1	94.8	93.8	93.2	1.06	0.586	0.467	0.506
TTTD of AEE, %		93.6	91.5	93.2	93.7	92.0	93.9	1.31	0.587	0.108	0.959

613 pSEM, Pooled SEM

614 ^{a,b,c}Means within a row lacking a common superscript letter are different ($P \leq 0.05$).

615 ¹ Least squares means of 9 pigs per diet.

616 ²SBO = soybean oil; DDGS = reduced oil distillers dried grains with solubles.






Fig. 1. Response surface for the effects of the dietary soybean oil (SBO) and reduced oil distillers dried grains with solubles (DDGS) on the apparent ileal (AID; A) and total tract digestibility (ATTD; B) of NDF in growing pigs ($n = 9$).



SBO × DDGS

$P = 0.004$

$R^2 = 0.79$





Fig. 2.

Response surface for the effects of the dietary soybean oil (SBO) and reduced oil distillers dried grains with solubles (DDGS) on the apparent ileal (AID; A) and total tract digestibility (ATTD; B) of acid hydrolyzed ether extract (AEE) in growing pigs ($n = 9$).



689

690 **Fig. 3.** Estimation of endogenous losses of acid hydrolyzed ether extract (AEE) by regression of
 691 total apparently digested AEE at the end of the ileum (TAD_i ; A) and over the entire intestinal
 692 tract (TAD_t ; B) as a function of dietary AEE intake. The y-intercepts are considered the
 693 estimated endogenous losses of AEE (g/kg of DMI) and the slope the true digestibility of AEE
 694 (Jørgensen et al., 1993).

695

696